
Implementation of Indiana's Probabilistic Watershed Monitoring Program and Second Year Source Identification Sampling





Presented by:

Stacey L. Sobat
Biological Studies Section
IDEM/Office of Water Quality/Assessment Branch

Overview of Presentation

- •Monitoring Design: why it was selected, how it is used, and products produced
- Second year source identification sampling
- •Strengths and weaknesses of using probabilistic monitoring





Goal of Indiana's Water Quality Standards

- •State's Goal: "... to restore and maintain the chemical, physical, and biological integrity of the waters of the State." 327 IAC 2-1-1.5
- ·Historically, targeted sites related to point source pollution
- •Shift to non-point source pollution & ALL waters of the state



Probabilistic Monitoring Design

•Rivers and streams are sampled by targeting watershed basins and sampling them every 5 years


-Year 1: WFWR & Patoka

-Year 2: EFWR & Whitewater

-Year 3: Upper Wabash


-Year 4: Lower Wabash & Kankakee

-Year 5: Great Lake & Ohio R. tribs



Probabilistic Monitoring Design

- •USEPA Western Ecology Division generated probabilistic site locations
- •Site Reconnaissance (Feb.-April)
- -Topographic Maps
- -Brochures
- -Contact Landowner
- -Site Accessibility & Equipment
- -Enter Recon data
- •Data Collected:
- -Water, Nutrient, and Bacteriological Samples For Laboratory Analysis
- -Macroinvertebrate Community Assessments
- -Fish Community Assessments
- -Habitat Assessments



Water Chemistry, Nutrients, & E.coli

- •Grab water samples and nutrients collected spring, summer, and fall (USGS contract)
- Chemistry Parameters

Priority Metals	Physical/Anions	Nutrients/Organic
Arsenic	Alkalinity	TKN
Calcium	Total Solids	Ammonia-N
Cadmium	Suspended Solids	Nitrate+Nitrite-N
Chromium	Dissolved Solids	Total Phosphorus
Copper	Sulfate	TOC
Lead	Chloride	Cyanide-Total
Magnesium	Hardness	Cyanide-Free
Mercury		COD
Nickel		
Selenium		
Zinc		

Water Chemistry, Nutrients, & E.coli

- •Hydrolab data collected during each sampling event:
- -D.O., pH, Conductivity, Water temperature, Turbidity
- •Nutrients:
- -Chlorophyll A in phytoplankton and periphyton samples, ash-free dry mass in periphyton samples
- •E.coli:
- —each site sampled once each week for 5 consecutive weeks





Macroinvertebrate Community Assessments

- •KICK sample
- •Hester-Dendy
- Laboratory Processing
- •mIBI: 10 metrics (range = 0-8)
- **−1.** Family level HBI
- **−2.** Number of taxa
- -3. Number of individuals
- -4. Percent dominant taxa
- -5. EPT Index
- -6. EPT Count
- -7. EPT count to total number of individuals
- **–8.** EPT Count to Chironomid Count
- -9. Chironomid Count
- -10. Total number of individuals to number of squares sorted





Fish Community Assessments

- •15 x the wetted stream width (backpack, totebarge, boat electrofishing)
- All stream sizes included
- Species Level Identification
- •IBI: 12 metrics (range 6-60)
- −1. # species
- -2. # darter species/ #dms species
- -3. % headwater ind./ # sunfish sp.
- -4. # minnow sp./ # sucker sp.
- **−5.** # sensitive sp.
- -6. % tolerant ind.
- −7. % omnivore ind.
- -8. % insectivore ind.
- -9. % pioneer ind./% carnivore ind.
- -10. Total # ind.
- **−11.** % simple lithophils
- **−12.** % DELT anomalies





Qualitative Habitat Evaluations (QHEI)

- 1. Substrate
- 2. Instream Cover
- 3. Channel Morphology
- 4. Riparian Zone & Bank Erosion
- 5. Pool/Glide Quality Riffle/Run Quality
- Gradient

Range 0-100

	73
EGICHAL TO THE	Qualitative Habitat Evaluation Index Field Sheet QHEI Score: 13
Event I.D. 00016	Stream West Branch Mosquito Creek County Harrison
Date 7-25-00 Loc	ation Ct 111000 Ct/15
- 51 AA	Crow Print City Cold Index Miles
11 SUBSTRATE (Check C	ONLYTwo Substrate TYPE BOXES; Estimate % present;
TYPE POOL	PIFFLE POOL HIFFLE SUBSTRATE UNION A LIVERACE COMMITTEE ON THE COMMITTEE O
J J-BLDR /SLBS[10]	
O O-BOULDER (9)	- CUT LODGE TE I II Substrate
□ 51-COBBLE [8] ✓	The state of the s
O O-HARDPAN (4)	Surface in
□ □-MUCK [2]	- G-ARTIFICIAL[0] - G-HARDPAN (0) - SILT FREE [1] - G-SILT FREE [1] - G-SANDSTONE [0] EMBEDDED G-EXTENSIVE [-2] Max 20
□ □-SILT [2]	T LIGHT TE (11
NOTE : (Ignore sludge origina	TO MANAGEMENT OF THE PROPERTY
scare on natural substrates)	J-3 of More (2)
NUMBER OF SUBSTRATE	TYPES: 10-4 or Less [0]
COMMENTS	AMOUNT: (Check ONLY One or Cover
2] INSTREAM COVER	to the statement of the control of t
	TYPE (Check All That Apply) O-DEEPPOOLS > 70 cm [2] O-OXBOWS[1] O-EXTENSIVE > 75% [11]
DUNDERCUTBANKS[1]	
D-OVERHANGING VEGETA	(ATTEN II) DEPOSITION
TA-SHALLOWS (INSLOWW	O-NEARLYABSENT < 5%[I]
D-ROOTMATS[1] CC	OLOCY- (Check ONLY One PER Category OH check 2 and AVENAGE)
	ME OPMENT CHANNELLAICH STABILITY
MI-TO-MAN-	EXCELLENTIFI MUNONEIGI BLHIGHISI DISNAGGING DIMPOUND.
7-1	COCOUNT D-RECOVERED 41 D-MODERATE 2] D-RELOCATION D-ISLANDS D-ISLANDS
	CLRECOVERING(3) O-LOWIII O-CANOPIREMOVAL D-CETELD MAX 20
	-PAIR]
COMMENTS:	DECOVERY[1] DECOVERSIDE OF AN INCLUDING THE STATE OF THE
RIPARIAN WIDTH	FLOOD FEATIN GOADY (FAST 1887 1887 1887 1887 1887 1887 1887 188
L R (Per Bank)	L R (Most Predominant Per Bank) L R L R (Per Bank) L R (Nost Predominant Per Bank) L R (Per Bank) D CONSERVATIONTILIAGE[1] X X NONE/LITLE[3]
文(A:-WIDE>50m[4]	D STOREST, STANKE [5]
O-MODERATE 10-50n	n [3] D D SHKUBOKOLD HED [2] B D O O O O O O O O O O O O O O O O O O
III-NARROWS-10m/2	Distance in the control limit
O O - VERYNARROW <	m[1] O O-FENCED PASTURE [1] O OMINING/CONSTRUCTION[U]
00NONE[0]	
COMMENTS:	THE OWN TV
5.]POOL/GLIDE AND	RIFFLE/RUN QUALITY Pool/
	MORPHOLOGY CURRENT VELOCITY [POOL & RIFFLESI] Glide
MAX. DEPTH	(Check 1 or 2 & AVERAGE) (Check All That Apply)
(Check 1 ONLY!)	DOOL WIDTH > RIFFLE WIDTH [2] O'-EDDIES[1] O'-TORRENTIAL[-1]
□ - > lm [6]	D-POOLWIDTH=RIFFLEWIDTH[1] D'-FAST[1] D'-INTERSTITIAL[-1] Max 12
□-0.7-1m[4]	O -POOLWIDTH < RIFFLEW.[0] O -MODERATE[1] O -INTERMITTENT[-2]
21-0.4-0.7m[2]	A: STOM[1]
0-0.2-0.4m[1]	
□ - < 0.2m [POOL=	Riffle/Run
	CHECK ONE OR CHECK 2 AND AVERAGE
RIFFLE /RUN DEPTH	RIFFLE/RUN SUBSTRATE RIFFLE/RUN EMBEDDEDNESS 9
GENERALLY >1	0 cm:MAX > 50 [4] D-STABLE (e.g., Connie, Double) (2)
O - GENERALLY > 1	O cm: MAX < 50[3] M-MOD. STABLE (e.g., Large Grovel) [1] D-LOW [1]
GENERALLY 5-	10 cm/11
D-GENERALLY < 5 cm	o (RIFFI F= 0)
COMMENTS:	J. NORTHE Presidency
	Max 10
an animum (in)	TWO I DRAINAGE AREA (sq.mi.): 12
6] GRADIENT (ft/	mi): 24-4 DRAINAGE AREA (sq.mi.): 12 %POOL: 60 %GUDE: 96RUN: 0

1. Any violations chemical or biological

- -For chemical impairment: Water Quality Standards [327 IAC 2-1-6]
- -For biological impairment:

"all waters, except those designated as limited use, will be capable of supporting a well-balanced, warm water aquatic community."

[327 IAC 2-1-3(2)]

"well-balanced aquatic community" is "an aquatic community which is diverse in species composition, contains several different trophic levels, and is not composed mainly of strictly pollution tolerant species"

[327 IAC 2-1-9(49)].




Chironomid Photo by Dale Parker, AquaTax Consulting



Central stoneroller, Campostoma anomalum

Total IBI Score	Integrity Class	Attributes
58-60	Excellent	Comparable to best condition present in ecoregion conditions,
48-52	Good	exceptional assemblage of species. Decreased species richness (intolerant species in particular), sensitive species present. Deviation Minor.
40-44	Fair	Intolerant and sensitive species absent, skewed trophic structure. Deviation Slight.
28-34	Poor	Top carnivores and many expected species absent or rare, omnivores and tolerant species dominant. Deviation Moderate.
12-22	Very Poor	Few species and individuals present, tolerant species dominant, diseased fish frequent. Deviation severe.
<12	No Fish	No fish captured during sampling. Deviation very severe.



Photos by Dale Parker, AquaTax Consulting

1. Any violations chemical or biological:

- -Fish Community: Impaired IBI \leq 35
- -Macroinvertebrate:

Impaired KICK mIBI < 2.2

Impaired Hester-Dendy < 1.4

2. Look for possible cause/source:

-habitat, type of chemical violation, likely source

3. How far to apply impairment:

-tributary influence, land use characterization, confined feeding operations, permitted facilities





- 1. Any violations chemical or biological
- 2. Look for possible cause/source
- 3. How far to apply impairment
- 4. Refer impaired sites for Source ID
- 5. Assess sites for Indiana's Integrated Water Monitoring and Assessment Report
- 6. Predict % Miles Attaining
 Aquatic Life Use
 Technical Reports
 Data Requests

INDIANA INTEGRATED WATER MONITORING AND ASSESSMENT REPORT 2002

Section 305(b) Water Quality Report and Consolidated List Including Section 303(d) List of Impaired Waters (Outagery 5)



Indiana Department of Environmental Management Office of Water Quality Planning and Restoration Breach Indianapolis, Indiana




- 1. "R" **software:** free software developed by the R Development Core Team along with many other contributors to allow the exchange of statistical packages and programming knowledge to anyone with an internet browser (http://cran.us.r-project.org/).
- 2. Table compatible with "R" software
- 3. Example of "R" screen & language
- 4. Adjust weight function
- 5. Output from "R"
- -Basic Statistics (mean, variance, std.deviation)
- -Cumulative distribution of stream miles for a numeric value (i.e. IBI score)
- -Percent of stream miles attaining with confidence levels
- 6. Assessment for River Basin



<u>F</u> ile <u>E</u> dit	t <u>S</u> earch <u>H</u> o	elp													
	SiteName		County	RF3RCHID	1	MAP24K	MAP100K	MAP250K	Long.dd	Lat.dd	Stratum	MDCaty	Weight	Strahler	Order 🔺
OECO.96	SiteIDF	Basin	HUC14Co	de	Strahle	rOrder1	StatusB	Fish.Co	mmunity	Fish.IB	[Macroinu			
Macro.		Habitat				Combined			T.E						
INRB 01-	-201	PATOKA I	R	GIBSON	5120209	32.	.70	Union	PRINCETO	DN	VINCENNE	2	87.38542	<u>.</u>	
38.3771	18	6	12	31.47142	9			^r River	Lowland	INRB 01-2	201	Patoka	5.12E+12	<u>.</u>	5
TS	I	18	NS	NA	I	36	I	I	A						
INRB 01-	-202	PATOKA I	R	DUBOIS	5120209	10 22.	.53	Cuzco	JASPER	VINCENNE	2	86.72151	l	38.44116	
6	12	31.47143	29	3	Interio	r Plateau	ı	INRB 01-	202	Patoka	5.12E+12	2	3	TS	NI 💻
42	NS	NA	NI		NI	ΝI	A								
INRB 01-	-203		PIKE	5120209	356 0	.00	Otwell	PRINCET	ON	VINCENNE	2	87.10332	2	38.44934	
6	11	44.06	1	Interior	River	Lowland	INRB 01-2	203	Patoka	5.12E+12	2	1	NT	NS	NA
NS	NA	NS	NA	NS	NS	A									
INRB 01-		Youngs (43 0.		Greenbr		JASPER			86.52804		
38.4444		6	13		2				INRB 01-2	204	Patoka	5.12E+12	2	2	NT
NS	NA	NS	NA		NA	NS	NS	A							
INRB01-			GIBSON						ON			87.45415		38.41468	
6	13	22.03	2				INRB 01-2	205	Patoka	5.12E+12	2	2	TS	I	22
NS	NA	I	44		I	Α							_		
INRB 01-		PATOKA I		DOBOIS	5120209	10 8. r Plateau	. 00		JASPER			86.85355		38.44801	
6	12	31.47143		4	Interior	r Plateau	1	INRB 01-	206	Patoka	5.12E+12	2	4	TS	I
22	NS	NA	NI	54	1	1	A								
INRB 01-		Cup Cred				18 0.	. 66	verbeu	PRINCETO	NN	VINCENNE		87.12247		т.
38.3271		6	13		2				INRB 01-2	207	Ратока	5.12E+12	!	2	TS
I	30	NI	4 NUDOTO	_		_	I	A	IACDED	HTHOCHU	- 6	04 00077	,	00 05460	
INRB 01-			DUBOIS	5120209			HUNCING	ourg	JASPER	OTHCENNE	: 2	86.88277		38.35649	
6 NI	13 5.2	22.03	2 42	Interior NI	NI NI	u A	INRB 01-2	2 08	Расока	5.12E+12	2	2	12	ΝI	38
INRB 01-		I Honey Ci			N1 5120209		00	Oalil and	City	DDINCETO	161	VINCENNE		87.27051	
38.2752		6	геек 11		1				INRB01-2			5.12E+12			TS
NI	46	NS	NA		57		I		TIME 0 1-2	2 09	гасика	J. 12E+12	-	•	13
INRB 01-			DUBOIS	5120209	98				JASPER	HINCENNE	· c	86.78765		38.36598	
6	13	22.03	2	Interior			INRB 01-2			5.12E+12)	2			40
NI	5.2	ZZ.03	47		NI	A	TIMD 91 - 2	210	racuka	J. 12L* 12	-	2	13	111	40
INRB 01-		PATOKA I					2 h	lle1nen	PRINCETO	nn	IITNCENNE	2	87.00374		
38.3607		6	 12	31.47142		5	Interio	River	Lowland	TNRR01-2)11	.s Patoka		-	5
LD.	NS	NA	NS		NS	NA	NS	NS	A	111111111111111111111111111111111111111		racona	J. 12L · 12	•	•
INRB 01-		••••	DUBOIS	5120209			Hunting			VINCENNE	2	86.98246		38.30035	
6	11	44.06	1			.ov Lowland	INRRO1-2	212		5.12E+12		1			16
NS	NA NA	I	33		I	A			· acona	5	•			-	. •
INRB 01-		-	PIKE	5120209			Petersb	ıra	PRINCETO	DN	VINCENNE	2	87.25887	,	
38.3839		6	12	31.47142					Lowland						4
NT	NS	NA	NS		NS	NA	NS	NS	A				· -	-	-
INRB 01-		PATOKA I				12 21.			JASPER	VINCENNE	2	86.47832	2	38.44104	
6	12	31.4714				r Plateau		INRB01-	214	Patoka					NI
46	NI	3	NI		NI	NI	A							_	
INRB 01-	-215	FLAT CR		5120209			Otwell	PRINCET	ON	VINCENNE	S	87.0772	38.40348	1	6
















- **1. "R" software:** free software developed by the R Development Core Team along with many other contributors to allow the exchange of statistical packages and programming knowledge to anyone with an internet browser (http://cran.us.r-project.org/).
- 2. Table compatible with "R" software
- 3. Example of "R" screen & language
- 4. Adjust weight function
- 5. Output from "R"
- -Basic Statistics (mean, variance, std.deviation)
- -Cumulative distribution of stream miles for a numeric value (i.e. IBI score)
- -Percent of stream miles attaining with confidence levels
- 6. Assessment for River Basin





- 1. "R" software: free software developed by the R Development Core Team along with many other contributors to allow the exchange of statistical packages and programming knowledge to anyone with an internet browser (http://cran.us.r-project.org/).
- 2. Table compatible with "R" software
- 3. Example of "R" screen & language
- 4. Adjust weight function
- 5. Output from "R"
- -Basic Statistics (mean, variance, std.deviation)
- -Cumulative distribution of stream miles for a numeric value (i.e. IBI score)
- -Percent of stream miles attaining with confidence levels
- 6. Assessment for River Basin



- **1. "R" software:** free software developed by the R Development Core Team along with many other contributors to allow the exchange of statistical packages and programming knowledge to anyone with an internet browser (http://cran.us.r-project.org/).
- 2. Table compatible with "R" software
- 3. Example of "R" screen & language
- 4. Adjust weight function
- 5. Output from "R"
- -Basic Statistics (mean, variance, std.deviation)
- -Cumulative distribution of stream miles for a numeric value (i.e. IBI score)
- -Percent of stream miles attaining with confidence levels
- 6. Assessment for River Basin







6. Assessment for River Basin

Project Name	Target Population	*% Attainment*	*% Non-Attainment*	Confidence Level	*Confidence Interval*
West Fork	05120201	69%	31%	95%	15%
White River	05120202		1st order=4		
n=36	05120203		2nd order=5		
1st=13			3rd order=2		
2nd=14			4th+ order=0		
3rd=4			n= 11/36 Impaired		
4th+=5					
Patoka River	05120209	29%	71%	95%	15%
n=26			1st order=4		
1st=5			2nd order=7		
2nd=11			3rd order=0		
3rd=3			4th+ order=7		
4th+=7			n=18/26 Impaired		
Target populatio	n = 8 digit HUC				

Target population = 8 digit HUC

[%] Attainment = Combined Assessment of Biology (IBI>35, mIBI>2.2,HD>1.4) and No Chemical Parameter Violations % Non-Attainment = Combined Assessment of Biology (IBI<35, mIBI<2.2,HD<1.4) and Chemical Parameter Violations The Confidence Interval is the % Non-Attainment +/- the value for 95% Confidence Level.

^{*} These are values produced by IDEM staff using "R" (http://cran.us.r-project.org/) and commands provided by USEPA National Health and Environmental Effects Research Laboratory, Corvallis, Oregon.

Second Year Source Identification Sampling

Original Intent of Surface Water Quality Monitoring Strategy

- 1. Find Impairments
- 2. Return to impairments the next year to find source
- 3. Assessments for Integrated Water Monitoring & Assessment Report

Reality: Limited success implementing second year studies

2004: Second Year Source Identification Fully Implemented

- USGS collecting water chemistry and nutrient data at probabilistic sampling sites
- Shift field season schedule for biological sampling to June rather than July
- Prioritize watershed source identification studies
- Shifting and hiring of staff to form new crews targeting impaired watersheds



Second Year Source Identification Sampling

•Extensive Chemical Sampling

(impaired sites due to pollutants or point sources)

Extensive Biological & Chemical Sampling

(impaired sites due to biological impairment)

- -Background research of potential sources
- -Approximately 30-40 sites per watershed (4-6 watersheds/summer) to determine cause and source of biological impairment

Products

- -Know extent of impairment and source for stream segment assessments
- -Proper listing of the waterbody in Categories 1-5 on the 303(d) List of Impaired Waterbodies
- -Data for TMDL development if needed and watershed restoration plans
- -Referrals to Inspectors, County Sanitation Districts and Department of Health
- -Potential Water Quality Criteria Revisions

Advantages & Disadvantages of Probabilistic Sampling

•Advantages:

- -100% waters of the state assessed
- -monitoring long term watershed trends
- -discovering non-point source pollution
- —finding impairments at distant remote sites
- —can focus resources to watershed specific impairments
- -Bonus: biological expansion of species distribution, Threatened & Endangered Species, educate public and landowners in watershed

•Disadvantages:

- -Time, Access, Safety
- -Cause and source of impairment
- -Basins sampled only once every 5 years
- -Where are the other impairments?



Photo Credit: Rob Criswell
Bluebreast Darter, *Etheostoma camurum*



Conclusions

Probability Monitoring

- -Completed one cycle, after 2005 2nd cycle will be completed!
- -Assess 100% of the waters of the state for the Integrated Water Monitoring and Assessment Report
- •Follow up impairments with second year source id
- -intensive surveys both biological and chemical in impaired watersheds
- -evaluate tiered aquatic life uses for modified streams
- •Need:
- -Numeric biocriteria
- -Refined-designated uses
- -Model to predict where impairments might occur
- -Determine cause and source for Impaired Biotic Communities already listed

Acknowledgements

- •Dennis Clark, Assessment Branch Chief
- •C. Lee Bridges, Biological Studies Section Chief
- •Arthur Garceau, Surveys Section Chief
- •Tony Olsen, USEPA NHEERL Corvallis, Oregon
- Assessment Branch Staff

